

IncyteCARES Patient Assistance Program ENROLLMENT FORM



COMPLETE AND FAX ALL PAGES TO **1-877-801-3840**.

**For patients who are uninsured or have Medicare Part D and cannot afford their out-of-pocket costs.
Commercially insured patients are not eligible for this Program.**

1. PATIENT INFORMATION

First Name _____ MI _____ Last Name _____
Address _____ City _____ State _____ ZIP _____
Patient Date of Birth ____/____/____ Gender _____
Caregiver Name _____ Relationship to Patient _____

Patient Communications

Phone _____ Email (Required for e-signature) _____

2. PATIENT INSURANCE INFORMATION

☐ **CHECK IF PATIENT DOES NOT HAVE PRESCRIPTION INSURANCE** (Move to section 3)

Primary Insurance

Primary Insurance Name _____ Primary Insurance Phone _____
Policyholder Name _____ Policyholder Date of Birth _____
Policy ID Number _____ Group Number _____

Prescription (Rx) Insurance

Rx Insurance Name _____ Rx Policy ID Number _____ Rx Group ID Number _____ Rx BIN _____ Rx PCN _____

3. PRESCRIBER INFORMATION

First Name _____ Last Name _____
Practice Name _____ State License Number _____ NPI Number _____
Address _____ City _____ State _____ ZIP _____
Practice Contact Name _____ Phone _____ Fax _____

4. PRESCRIPTION FOR OPZELURA

Primary Diagnosis ICD-10 Code(s) ☐ L20 Atopic Dermatitis ☐ L80 Vitiligo ☐ Other ICD-10 _____

Medication Name: **OPZELURA® (ruxolitinib) cream 1.5%** Tube size: 60 grams Number of Tubes _____ Refill(s) _____
Directions _____

I certify that I am the Healthcare Professional who has prescribed this medication, that it is medically necessary for the patient, and that the information provided is accurate to the best of my knowledge. I authorize Incyte, and its affiliates, agents, and service providers to act on my behalf for the purposes of transmitting this prescription to the appropriate pharmacy.

If you are a prescriber based in New York state, please use a New York state prescription form.

**PRESCRIBER
SIGNATURE** X

Dispense as written

X

Substitutions allowed

____/____/____

Date

5. PRESCRIBER DECLARATION

By my signature, I certify that I have obtained any and all authorizations and consents from the patient or the patient's authorized personal representative necessary under HIPAA and state law to release protected health information, including that contained on this form, to Incyte and its employees or agents for purposes relating to Incyte's patient support programs.

**PRESCRIBER
SIGNATURE** X

____/____/____

Date

IncyteCARES Patient Assistance Program ENROLLMENT FORM



COMPLETE AND FAX ALL PAGES TO **1-877-801-3840**.

Provide a copy of the HIPAA authorization to your patient for their records.

HIPAA AUTHORIZATION

By signing this form, you are giving your permission to your physicians, pharmacies, and other healthcare providers (collectively, "Healthcare Professionals"), your health insurer, health plan or programs that provide your healthcare benefits (collectively, "Health Insurers"), to share your health information with Incyte, its agents, and Incyte patient support program service providers (collectively, "Incyte"). You understand that your health information includes information relating to your medical condition and treatment, health insurance coverage and claims, and your prescription (including fill/refill information) for OPZELURA, as well as identifying information about you (including, for example, your name, address, and date of birth) ("Your Information"). Your information will be shared with Incyte so that Incyte may provide you with various support and information to help you access OPZELURA, which may include the following (collectively, "Patient Support Programs"):

- Providing reimbursement support, including assisting with identification of your insurer's prior authorization requirements and requirements for appealing a denied claim
- Provide adherence reminders and support
- Determining your eligibility for and helping you access copay support for OPZELURA
- Determining your eligibility to participate in coverage assistance programs, patient assistance programs, or other support programs
- Communicating with your Healthcare Professionals about OPZELURA and available Patient Support Programs
- Providing you with financial assistance resources and information if you are eligible
- Communication from Incyte regarding the Patient Support Programs using phone, email, text, or an auto dialer or prerecorded voice using the information provided to enable fulfillment of the Patient Support Programs described above.

You understand that you do not have to sign this form and choosing not to sign will not affect your ability to receive treatment from your Healthcare Professionals or payment from your Health Insurer.

However, if you do not sign this form, Incyte will not be able to provide you with assistance.

You understand that once Your Information is shared, it may no longer be protected by federal privacy law. However, Incyte agrees to protect Your Information and to use it only for the purposes described in this form or as required or permitted by law.

You understand that this form will remain in effect for two (2) years from the date of your signature unless you provide written notice that you would like to withdraw your authorization to share your health information sooner.

If you would like to withdraw your authorization, you may contact IncyteCARES for OPZELURA at 1-800-583-6964 or PO Box 7613, Overland Park, KS 66207. You understand that if you withdraw your authorization, no new information will be collected from you; however, Your Information collected prior to your withdrawal of authorization may continue to be used or kept to provide services previously described. You understand you may receive a copy of this form.

Incyte also may use Your Information for quality assurance purposes and to evaluate and improve its operations and services. You also understand that the information you provide may be combined with that of other registrants to create aggregated, anonymized data and to use and share only the anonymized data for any legitimate business purpose. You can learn more about how Incyte processes Your Information at www.incyte.com/privacy-policy.

PATIENT
SIGNATURE **X**

(If the patient is under 18 years of age, a legal representative should sign and print their name.)

Date

Legal Representative Name (Print) _____

IncyteCARES Patient Assistance Program ENROLLMENT FORM



COMPLETE AND FAX ALL PAGES TO **1-877-801-3840**.

Patient First Name _____ MI _____ Patient Last Name _____ Date of Birth ____/____/____

AUTHORIZATION FOR ELECTRONIC INCOME VERIFICATION

I, the applicant named above, understand that I am providing "written instructions" to Incyte under the Fair Credit Reporting Act authorizing Incyte or their designated agent to obtain information from my credit profile or other information from TransUnion to verify my annual income. I authorize Incyte to obtain such information solely for the purpose of determining financial qualifications for the Patient Assistance Program. I also agree to provide additional financial documentation in a timely manner to Incyte, which is solely necessary for determining my financial qualification for the program, if so requested. I understand that I am entitled to a copy of this authorization upon request. This authorization shall be valid for one (1) year from the date of the signature on this form (unless a shorter period is prescribed by law). I understand that I may cancel this authorization at any time by mailing a letter requesting such cancellation to Incyte at PO Box 7613, Overland Park, KS 66207. I understand that if I cancel my authorization, no new information will be collected from me; however, information collected prior to such cancellation may continue to be used or kept to fulfill the services described.

Please note: If you do not provide an authorization signature, you are required to submit income verification via fax to 1-877-801-3840, or mail to Incyte at PO Box 7613, Overland Park, KS 66207.

PATIENT
SIGNATURE

X

(If the patient is under 18 years of age, a legal representative should sign and print their name.)

____/____/____
Date

Legal Representative Name (Print) _____

FINANCIAL ATTESTATION AND DISCLOSURE

By signing below, I certify that I cannot afford my medication, and I affirm that my answers are complete, true, and accurate to the best of my knowledge.

I understand that: Completing this form does not guarantee that I will qualify for the Patient Assistance Program. Incyte may verify the accuracy of the information I have provided and may ask for more financial and insurance information. OPZELURA supplied by the Patient Assistance Program shall not be sold, traded, bartered, or transferred. Incyte reserves the right to change its Patient Assistance Program or terminate my enrollment at any time. The support provided through this program is not contingent on any future purchase. If I am enrolled in a Medicare Part D Plan and am eligible for the Patient Assistance Program, Incyte will notify my Part D Plan of my enrollment in the program.

I certify and attest that if I receive OPZELURA provided through the Incyte Patient Assistance Program, I will promptly contact the program at 1-800-583-6964 if my financial status or insurance coverage changes. I will not seek to have OPZELURA or any cost from it counted in my Medicare Part D true out-of-pocket costs for prescription drugs. I will not seek reimbursement or credit for OPZELURA from my prescription insurance provider or payer, including Medicare Part D plans. I will notify my insurance provider of the receipt of OPZELURA through the Incyte Patient Assistance Program.

For more information about Incyte and its privacy practices, please go to www.incyte.com/privacy-policy.

PATIENT
SIGNATURE

X

(If the patient is under 18 years of age, a legal representative should sign and print their name.)

____/____/____
Date

Legal Representative Name (Print) _____

PATIENT OPT-IN FOR ONGOING EDUCATION AND SUPPORT (OPTIONAL)

- ☐ By checking this box, I consent to Incyte and its agents to use my contact information (phone and email) to provide education and ongoing support services related to product, disease, and other related areas of interest. I understand that I may revoke my consent to be contacted for any of these purposes at any time by emailing privacy@incyte.com.
- ☐ By checking this box, I am indicating that I would also like to be contacted for future opportunities to participate in market research regarding my condition. I understand that I may revoke my consent to be contacted to participate in such market research at any time by emailing privacy@incyte.com.

X

Patient Signature (If the patient is under 18 years of age, a legal representative should sign and print their name.)

____/____/____
Date

Legal Representative Name (Print) _____

INDICATIONS

OPZELURA is indicated for the topical short-term and non-continuous chronic treatment of mild to moderate atopic dermatitis in non-immunocompromised adult and pediatric patients 12 years of age and older whose disease is not adequately controlled with topical prescription therapies or when those therapies are not advisable.

OPZELURA is indicated for the topical treatment of nonsegmental vitiligo in adult and pediatric patients 12 years of age and older.

Limitations of Use: Use of OPZELURA in combination with therapeutic biologics, other JAK inhibitors, or potent immunosuppressants such as azathioprine or cyclosporine is not recommended.

IMPORTANT SAFETY INFORMATION

SERIOUS INFECTIONS

Patients treated with oral Janus kinase inhibitors for inflammatory conditions are at risk for developing serious infections that may lead to hospitalization or death. Reported infections include:

- Active tuberculosis, which may present with pulmonary or extrapulmonary disease.
- Invasive fungal infections, including cryptococcosis and pneumocystosis.
- Bacterial, viral, including herpes zoster, and other infections due to opportunistic pathogens.

Avoid use of OPZELURA in patients with an active, serious infection, including localized infections. If a serious infection develops, interrupt OPZELURA until the infection is controlled. Carefully consider the benefits and risks of treatment prior to initiating OPZELURA in patients with chronic or recurrent infection. Closely monitor patients for the development of signs and symptoms of infection during and after treatment with OPZELURA.

Serious lower respiratory tract infections were reported in the clinical development program with topical ruxolitinib.

No cases of active tuberculosis (TB) were reported in clinical trials with OPZELURA. Cases of active TB were reported in clinical trials of oral Janus kinase inhibitors used to treat inflammatory conditions. Consider evaluating patients for latent and active TB infection prior to administration of OPZELURA. During OPZELURA use, monitor patients for the development of signs and symptoms of TB.

Viral reactivation, including cases of herpes virus reactivation (e.g., herpes zoster), were reported in clinical trials with Janus kinase inhibitors used to treat inflammatory conditions including OPZELURA. If a patient develops herpes zoster, consider interrupting OPZELURA treatment until the episode resolves.

Hepatitis B viral load (HBV-DNA titer) increases, with or without associated elevations in alanine aminotransferase and aspartate aminotransferase, have been

reported in patients with chronic HBV infections taking oral ruxolitinib. OPZELURA initiation is not recommended in patients with active hepatitis B or hepatitis C.

MORTALITY

In a large, randomized, postmarketing safety study in rheumatoid arthritis (RA) patients 50 years of age and older with at least one cardiovascular risk factor comparing an oral JAK inhibitor to tumor necrosis factor (TNF) blocker treatment, a higher rate of all-cause mortality, including sudden cardiovascular death, was observed with the JAK inhibitor.

Consider the benefits and risks for the individual patient prior to initiating or continuing therapy with OPZELURA.

MALIGNANCIES

Malignancies were reported in patients treated with OPZELURA. Lymphoma and other malignancies have been observed in patients receiving JAK inhibitors used to treat inflammatory conditions. In RA patients treated with an oral JAK inhibitor, a higher rate of malignancies (excluding non-melanoma skin cancer (NMSC)) was observed when compared with TNF blockers. Patients who are current or past smokers are at additional increased risk.

Consider the benefits and risks for the individual patient prior to initiating or continuing therapy with OPZELURA, particularly in patients with a known malignancy (other than successfully treated non-melanoma skin cancers), patients who develop a malignancy when on treatment, and patients who are current or past smokers.

Non-melanoma skin cancers, including basal cell and squamous cell carcinoma, have occurred in patients treated with OPZELURA. Perform periodic skin examinations during OPZELURA treatment and following treatment as appropriate. Exposure to sunlight and UV light should be limited by wearing protective clothing and using broad-spectrum sunscreen.

MAJOR ADVERSE CARDIOVASCULAR EVENTS (MACE)

In RA patients 50 years of age and older with at least one cardiovascular risk factor treated with an oral JAK inhibitor, a higher rate of major adverse cardiovascular events (MACE) (defined as cardiovascular death, myocardial infarction, and stroke), was observed when compared with TNF blockers. Patients who are current or past smokers are at additional increased risk. Discontinue OPZELURA in patients who have experienced a myocardial infarction or stroke.

Consider the benefits and risks for the individual patient prior to initiating or continuing therapy with OPZELURA, particularly in patients who are current or past smokers and patients with other cardiovascular risk factors. Patients should be informed about the symptoms of serious cardiovascular events and the steps to take if they occur. Discontinue OPZELURA in patients that have experienced a myocardial infarction or stroke.

THROMBOSIS

Thromboembolic events were observed in trials with OPZELURA. Thrombosis, including pulmonary embolism (PE), deep venous thrombosis (DVT), and arterial thrombosis have been reported in patients receiving JAK inhibitors used to treat inflammatory conditions. Many of these adverse reactions were serious and some resulted in death. In RA patients 50 years of age and older with at least one cardiovascular risk factor treated with an oral JAK inhibitor, a higher rate of thrombosis was observed when compared with TNF blockers. Avoid OPZELURA in patients at risk. If symptoms of thrombosis occur, discontinue OPZELURA and treat appropriately.

Thrombocytopenia, Anemia, and Neutropenia

Thrombocytopenia, anemia, and neutropenia were reported in the clinical trials with OPZELURA. Consider the benefits and risks for individual patients who have a known history of these events prior to initiating therapy with OPZELURA. Perform CBC monitoring as clinically indicated. If signs and/or symptoms of clinically significant thrombocytopenia, anemia, and neutropenia occur, patients should discontinue OPZELURA.

Lipid Elevations

Treatment with oral ruxolitinib has been associated with increases in lipid parameters including total cholesterol, low-density lipoprotein (LDL) cholesterol, and triglycerides.

Adverse Reactions

In atopic dermatitis, the most common adverse reactions ($\geq 1\%$) are nasopharyngitis (3%), diarrhea (1%), bronchitis (1%), ear infection (1%), eosinophil count increased (1%), urticaria (1%), folliculitis (1%), tonsillitis (1%), and rhinorrhea (1%).

In nonsegmental vitiligo, the most common adverse reactions (incidence $\geq 1\%$) are application site acne (6%), application site pruritus (5%), nasopharyngitis (4%), headache (4%), urinary tract infection (2%), application site erythema (2%), and pyrexia (1%).

Pregnancy Registry

There is a pregnancy registry that monitors pregnancy outcomes in pregnant persons exposed to OPZELURA during pregnancy. Pregnant persons exposed to OPZELURA and healthcare providers should report OPZELURA exposure by calling 1-855-463-3463 or visiting www.opzelura.pregnancy.incyte.com.

Lactation

Advise women not to breastfeed during treatment with OPZELURA and for approximately four weeks after the last dose (approximately 5-6 elimination half-lives).

Please see Full Prescribing Information, including Boxed Warning, and Medication Guide for OPZELURA.